

BOTANICAL ANTI-BACTERIAL BLEND

Hydrosoluble



ALOE VERA - LIME - NEEM - ROSEMARY - SAGE - THYME - VETIVER

The BOTANICAL ANTI-BACTERIAL BLEND is a multifunctional botanical complex selected for the complementary abilities of each single extract in order to prevent microorganisms proliferation on skin and hair.

The BOTANICAL ANTI-BACTERIAL BLEND combines leaves, fruit and root, which were selected for their synergetic and complementary activities:

- ✓ ***Antiseptic***
- ✓ ***Anti-fungal / Anti-dandruff***
- ✓ ***Insect repellent***
- ✓ ***Anti-dehydration / Brings moisturization / Refreshing***
- ✓ ***Soothing / Calming / Softening***
- ✓ ***Anti-ageing: - Antioxidant***
- Regenerating
- ✓ ***Circulatory / Detoxifying***
- ✓ ***Astringent***



COSMETIC USES

The **BOTANICAL ANTI-BACTERIAL BLEND** is an efficient active, by following effects:

- > **Antiseptic**
- > **Anti-fungal / Anti-dandruff**
- > **Insect repellent**
- > **Anti-dehydration / Brings moisturization / Refreshing**
- > **Soothing / Calming / Softening**
- > **Anti-ageing: Antioxidant / Regenerating**
- > **Circulatory / Detoxifying**
- > **Astringent**



INCI Name: Aloe Barbadensis Leaf Extract, Citrus Aurantifolia (Lime) Fruit Extract, Azadirachta Indica Leaf Extract, Rosmarinus Officinalis (Rosemary) Leaf Extract, Salvia Officinalis (Sage) Leaf Extract, Thymus Vulgaris (Thyme) Leaf Extract, Vetiveria Zizanioides Root Extract

CAS N°: 85507-69-3, 90063-52-8, - , 84604-14-8, 84082-79-1, 84929-51-1, 84238-29-9
EINECS N°: 287-390-8, 290-010-3, - , 283-291-9, 282-025-9, 284-535-7, 282-490-8

BIBLIOGRAPHIC REFERENCES

- ANSM. Pharmacopée française 2003.
- Barrau V. 2011. L'herbier d'une vie : Quand les plantes nous accompagnent du premier au dernier souffle. Plumes de carotte.
- Bruneton J. 2009. Pharmacognosie, phytochimie, Plantes médicinales. Editions Lavoisier - Tec&Doc.
- Demo, A., Petrakis, C., Kefalas, P., & Boskou, D. (1998). Nutrient antioxidants in some herbs and Mediterranean plant leaves. Food research international, 31(5), 351-354.
- Đilas, S., Knez, Ž., Četojević-Simin, D., Tumbas, V., Škerget, M., Čanadanović-Brunet, J., & Četković, G. (2012). In vitro antioxidant and antiproliferative activity of three rosemary (Rosmarinus officinalis L.) extract formulations. International Journal of Food Science & Technology, 47(10), 2052-2062.
- Dinda, A., Das, D., Ghosh, G., & Kumar, S. (2013). Analgesic and Anti-inflammatory Activity of Various Fractions of Azadirachta indica Leaf in Experimental Animals. International Journal of PharmTech Research, 5(2).
- Duke JA. 2002. Handbook of medicinal herb 2nd édition. CRC Press.
- Dweck AC. 2011. Formulating natural cosmetics. Allured books.
- Erkan, N., Ayranci, G., & Ayranci, E. (2008). Antioxidant activities of rosemary (R. Officinalis L.) extract, blackseed (N. sativa L.) essential oil, carnosic acid, rosmarinic acid and sesamol. Food Chemistry, 110(1), 76-82.
- Gupta, A., Malviya, R., Singh, T. P., & Sharma, P. K. (2010). Indian medicinal plants used in hair care cosmetics: a short review. Pharmacognosy Journal, 2(10), 361- 364.
- Hidalgo, P. J., Ubera, J. L., Tena, M. T., & Valcarcel, M. (1998). Determination of the carnosic acid content in wild and cultivated Rosmarinus officinalis. Journal of Agricultural and Food Chemistry, 46(7), 2624-2627.
- Justo, O. R., Simioni, P. U., Gabriel, D. L., Tamashiro, W. M., Rosa, P. T., & Moraes, Á. M. (2015). Evaluation of in vitro anti-inflammatory effects of crude ginger and rosemary extracts obtained through supercritical CO2 extraction on macrophage and tumor cell line. BMC complement altern med, 15(1), 390.
- Kuo, C. F., Su, J. D., Chiu, C. H., Peng, C. C., Chang, C. H., Sung, T. Y., ... & Chyau, C. C. (2011). Anti-inflammatory effects of supercritical carbon dioxide extract and its isolated carnosic acid from Rosmarinus officinalis leaves. Journal of agricultural and food chemistry, 59(8), 3674-3685.

- Mahabub-Uz-Zaman, M., Ahmed, N. U., Akter, R., Ahmed, K., Aziz, M. S. I., & Ahmed, M. S. (2009). Studies on anti-inflammatory, antinociceptive and antipyretic activities of ethanol extract of *Azadirachta indica* leaves. *Bangladesh Journal of Scientific and Industrial Research*, 44(2), 199-206.
- Martini MC and Seiller M. 2006. *Actifs et additifs en cosmetology*. Editions Lavoisier Tec&Doc.
- Mosaddek, A. S. M., & Rashid, M. M. U. (2008). A comparative study of the anti-inflammatory effect of aqueous extract of neem leaf and dexamethasone. *Bangladesh Journal of Pharmacology*, 3(1), 44-47.
- Natarajan, V., Venugopal, P. V., & Menon, T. (2003). Effect of *Azadirachta indica* (neem) on the growth pattern of dermatophytes. *Indian journal of medical microbiology*, 21(2), 98.
- Naveena, B. M., Vaithyanathan, S., Muthukumar, M., Sen, A. R., Kumar, Y. P., Kiran, M., ... & Chandran, K. R. (2013). Relationship between the solubility, dosage and antioxidant capacity of carnosic acid in raw and cooked ground buffalo meat patties and chicken patties. *Meat science*, 95(2), 195-202.
- Niharika, A., Aquicio, J. M., & Anand, A. (2010). Antifungal properties of neem (*Azadirachta indica*) leaves extract to treat hair dandruff. *E-International Scientific International Scientific International Scientific Research Journal Research Journal*, 244.
- Okpanyi, S. N., & Ezeukwu, G. C. (1981). Anti-Inflammatory and Antipyretic Activities of *Azadirachta indica**. *Planta medica*, 41(01), 34-39.
- Perez-Fons, L., Garzón, M. T., & Micol, V. (2009). Relationship between the antioxidant capacity and effect of rosemary (*R. officinalis* L.) polyphenols on membrane phospholipid order. *J Agric Food Chem*, 58(1), 161-171.
- Sharma, L., Agarwal, G., & Kumar, A. (2003). Medicinal plants for skin and hair care. *Indian Journal of Traditional Knowledge*, 2(1), 62-68.
- Singh, K. K. (2009). *Neem, a Treatise*. IK International Pvt Ltd.
- Rignar, K., Celan, S., Skerget, M., & Knez, Z. (2008). Solubility of carnosic acid and carnosol from rosemary extract in supercritical CO₂. *Acta chimica slovenica*, 55(3), 516-520.
- Senorans, F. J., Ibanez, E., Cavero, S., Tabera, J., & Reglero, G. (2000). Liquid chromatographic-mass spectrometric analysis of supercritical-fluid extracts of rosemary plants. *J of Chromatography A*, 870(1), 491-499.
- Tena, M. T., Valcárcel, M., Hidalgo, P. J., & Ubera, J. L. (1997). Supercritical fluid extraction of natural antioxidants from rosemary: comparison with liquid solvent sonication. *Analytical chemistry*, 69(3), 521-526.
- Troncoso, N., Sierra, H., Carvajal, L., Delpiano, P., & Günther, G. (2005). Fast high performance liquid chromatography and ultraviolet-visible quantification of principal phenolic antioxidants in fresh rosemary. *Journal of Chromatography A*, 1100(1), 20-25.
- Ulbricht, C., Abrams, T. R., Brigham, A., Ceurvels, J., Clubb, J., Curtiss, W., ... & Windsor, R. C. (2010). An evidence-based systematic review of rosemary (*Rosmarinus officinalis*) by the Natural Standard Research Collaboration. *Journal of dietary supplements*, 7(4), 351-413.
- <https://www.botanical.com>, <http://www.catalogueoflife.org>, <http://ec.europa.eu>, <https://hort.purdue.edu>, <http://online.personalcarecouncil.org>, <http://www.pfaf.org>, <http://sun.ars-grin.gov>.